



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\frac{\int \cdots \int_T dx_1 \cdots dx_n}{\int \cdots \int_S dx_1 \cdots dx_n};$$

in other words, find the magnitude of a "solid angle" in n dimensions, with the "sphere" as unit solid angle.

Note. This problem was discussed and left unsolved by Schläfli in the *Quarterly Journal of Mathematics* for 1858, 1860, 1867.—Editor.

349 (Calculus) [December, 1913]. Proposed by C. N. SCHMALL, New York City.

If $y = a \cos (\log x) + b \sin (\log x)$, eliminate the constants a and b and obtain the equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + y = 0.$$

198 (Number Theory) [November, 1913]. Proposed by the late ARTEMAS MARTIN.

Prove that every even number is the sum of two prime numbers.

Note. This problem has long been known and no proof has ever been given.—Editor.

201 (Number Theory) [December, 1913]. Proposed by E. T. BELL, University of Washington.

Eisenstein proposed (*Crelle*, t. 27, p. 282), as the simplest of several problems: "In the expansion of

$$\frac{1 + z + z^2 + \cdots + z^{p-1}}{(1 - z)^{p-1}} - 1,$$

where p is prime, to show that the coefficients of the various powers of z are all divisible by p ."

SOLUTIONS OF PROBLEMS.

231 (Number Theory) [April, 1915]. Proposed by A. J. KEMPNER, University of Illinois.

Is the series whose terms are the reciprocals of all positive integers not containing a given combination of figures, for example not containing the combination 37, convergent or divergent? Numbers such as $\frac{1}{37}$, $\frac{1}{370}$, $\frac{1}{371}$ shall be omitted, numbers such as $\frac{1}{73}$, $\frac{1}{307}$, $\frac{1}{5317}$ shall be admitted as terms of the series. (Compare AMERICAN MATHEMATICAL MONTHLY, Volume 21, page 123.)

SOLUTION BY FRANK IRWIN, University of California.

Let $f(n)$ be the number of numbers in the class, g_n , of numbers of n digits that do not contain the given combination, for example 37. We can get these numbers by taking any number of g_{n-1} and adding a digit at the end; except that this digit must not be 7 if the number chosen from g_{n-1} ends with a 3. Since the number of these rejected numbers of g_{n-1} is evidently $f(n-2)$, we have the formula: $f(n) = 10f(n-1) - f(n-2)$. In the general case, where the given combination consists of k digits, we should get similarly, for $n > k$,

$$f(n) = 10f(n-1) - f(n-k). \quad (1)$$

(An exception would arise for such a combination as 37537; here, for instance, the formula $f(8) = 10f(7) - f(3)$ would not hold, for the number 375 belonging to g_3 would have been subtracted, as it should not have been, since the number obtained by writing after it 3753, viz., 3753753, does not belong to g_7 , and so has not been counted in $f(7)$. It will be seen that such cases arise when the first l digits of the given combination are the same as the last l , $l < k$. We shall call such combinations "improper" and deal with them separately.)

From (1) we have $f(n-1) > f(n)/10$. Also since $f(n-1) = f(n)/10$, $n < k$, and $f(n-1) \geq f(n)/10$, $n = k$ (= if the given combination begins with a zero), we have, for $n > k$, the string of relations $f(n-2) \geq f(n-1)/10$, $f(n-3) \geq f(n-2)/10$, \dots . From these last inequalities we derive $f(n-k) \geq f(n-1)/10^{k-1}$. Comparing this with (1), we see that $f(n) \leq 10f(n-1) - f(n-1)/10^{k-1}$, or $f(n) \leq 10^k f(n-1)$, if we put $(10^k - 1)/10^k = r$. Since the reciprocal of any number in g_n is $\leq 1/10^{n-1}$, it follows that the sum of the reciprocals of the numbers in g_k , g_{k+1} , \dots are less than the corresponding terms of the series